

Some Remarks on the Best Approximation Rate of Certain Trigonometric Series

Songping Zhou¹ and Ruijun Le

ABSTRACT. The main object of the present paper is to give a complete result regarding the best approximation rate of certain trigonometric series in general complex valued continuous function space under a new condition which gives an essential generalization to O -regularly varying quasimonotonicity. An application is present in Section 3.

1991 Mathematics Subject Classification. 42A20 42A32

§1. Introduction

As generalizations to monotonicity condition, various quasimonotonicity conditions including so-called O -regularly varying quasimonotonicity are introduced and applied to convergence of trigonometric (Fourier) series. Interested readers could check references such as Nurcombe [3], and Xie and Zhou [8] for uniform convergence and Stanojevic [4, 5], and Xie and Zhou [7] for L^1 convergence.

Let $C_{2\pi}$ be the space of all real or complex valued continuous functions $f(x)$ of period 2π with norm

$$\|f\| = \max_{-\infty < x < \infty} |f(x)|,$$

and $E_n(f)$ the best approximation by trigonometric polynomials of degree n . Given a trigonometric series $\sum_{k=-\infty}^{\infty} c_k e^{ikx} := \lim_{n \rightarrow \infty} \sum_{k=-n}^n c_k e^{ikx}$, write

$$f(x) = \sum_{k=-\infty}^{\infty} c_k e^{ikx} =: \sum_{k=-\infty}^{\infty} \hat{f}(k) e^{ikx}$$

at any point x where the series converges. Denote its n th partial sum $S_n(f, x)$ by

$$\sum_{k=-n}^n \hat{f}(k) e^{ikx}.$$

For a sequence $\{c_n\}_{n=0}^{\infty}$, let

$$\Delta c_n = c_n - c_{n+1}.$$

A non-decreasing positive sequence $\{R(n)\}_{n=1}^{\infty}$ is said to be O -regularly varying if²

$$\limsup_{n \rightarrow \infty} \frac{R(2n)}{R(n)} < \infty.$$

¹Supported in part by National and Zhejiang Provincial Natural Science Foundations of China under grant numbers 1041001 and 101009 respectively.

²In some papers, this requirement is written as that for some $\lambda > 1$, $\limsup_{n \rightarrow \infty} R([\lambda n])/R(n) < \infty$. It is just a pure decoration.

A complex sequence $\{c_n\}_{n=0}^{\infty}$ is O -regularly varying quasimonotone in complex sense if for some $\theta_0 \in [0, \pi/2)$ and some O -regularly varying sequence $\{R(n)\}$ the sequence

$$\Delta \frac{c_n}{R(n)} \in K(\theta_0) := \{z : |\arg z| \leq \theta_0\}, \quad n = 1, 2, \dots.$$

Evidently, if $\{c_n\}$ is a real sequence, then the O -regularly varying quasimonotonicity becomes

$$\Delta \frac{c_n}{R(n)} \geq 0, \quad n = 1, 2, \dots,$$

which was used in many works to generalize the regularly varying quasimonotone condition and, in particular, the quasimonotone condition³.

In 1992, without proof, Belov [1] announced the following result:

Theorem B. *Let $\{a_n\}$ and $\{b_n\}$ be quasimonotone sequences satisfying $\sum_{n=0}^{\infty} a_n < \infty$ and $\lim_{n \rightarrow \infty} nb_n = 0$. Define*

$$f(x) = \sum_{n=0}^{\infty} a_n \cos nx, \quad g(x) = \sum_{n=1}^{\infty} b_n \sin nx.$$

Then $f, g \in C_{2\pi}$ have the following estimates:

$$E_n(f) \approx \max_{1 \leq k \leq n} ka_{n+k} + \sum_{k=2n+1}^{\infty} a_k,$$

and

$$E_n(g) \approx \max_{k \geq 1} kb_{n+k}.$$

It is essentially generalized recently by Xiao, Xie and Zhou [6] for general complex valued trigonometric series (thus a proof is given to Theorem B) as follows.

Theorem XX. *Let $\{\hat{f}(n)\}_{n=0}^{\infty}$ and $\{\hat{f}(n) + \hat{f}(-n)\}_{n=1}^{\infty}$ be complex O -regularly varying quasimonotone sequences, and*

$$f(x) = \sum_{n=-\infty}^{\infty} \hat{f}(n) e^{inx}.$$

Then $f \in C_{2\pi}$ if and only if

$$\sum_{n=1}^{\infty} |\hat{f}(n) + \hat{f}(-n)| < \infty$$

and

$$\lim_{n \rightarrow \infty} n\hat{f}(n) = 0.$$

³A real sequence $\{b_n\}_{n=0}^{\infty}$ is defined to be quasimonotone if, for some $\alpha \geq 0$, the sequence $\{b_n/n^\alpha\}$ is non-increasing.

Furthermore, if $f \in C_{2\pi}$, then⁴

$$\begin{aligned} E_n(f) \approx & \max_{1 \leq k \leq n} k \left(\left| \hat{f}(n+k) \right| + \left| \hat{f}(-n-k) \right| \right) + \max_{k \geq 2n+1} k \left| \hat{f}(k) - \hat{f}(-k) \right| \\ & + \sum_{k=2n+1}^{\infty} |\hat{f}(k) + \hat{f}(-k)|. \end{aligned}$$

We make a quick remark here. We notice that, in the complex valued function space, the assumption that both $\{\hat{f}(n)\}_{n=0}^{\infty}$ and $\{\hat{f}(-n)\}_{n=1}^{\infty}$ are complex O -regularly varying quasimonotone is a convenient one, but is almost trivial, since it is almost the same as the condition in real case. Thus people usually use one side quasimonotonicity with some kind balance condition in considering those problems. Theorem XZ reflects this kind of thinking.

Motivated by an idea in Leindler [2], our work [11] introduced a new condition, and we present here a revised form as follows:

Definition. Let $\mathbf{c} = \{c_n\}_{n=1}^{\infty}$ be a sequence satisfying $c_n \in K(\theta_1)$ for some $\theta_1 \in [0, 2\pi)$ and $n = 1, 2, \dots$. If there is a natural number N_0 such that

$$\sum_{n=m}^{2m} |\Delta c_n| \leq M(\mathbf{c}) \max_{m \leq n < m+N_0} |c_n| \quad (1)$$

holds for all $m = 1, 2, \dots$, where $M(\mathbf{c})$ indicates a positive constant only depending upon $\mathbf{c}$, then we say that the sequence $\mathbf{c}$ belongs to class GBV.

We recall the following results.

Lemma 1. Suppose a complex sequence $\{c_n\}$ is O -regularly varying quasimonotone, then there is a positive constant M depending upon θ_0 only such that

$$|c_n| \leq M \text{Rec}_n, \quad n = 1, 2, \dots,$$

or in other words, $c_n \in K(\theta_1)$ for some $\theta_1 \in [0, \pi/2)$ and $n = 1, 2, \dots$.

The argument exactly follows from Xie and Zhou [8, Lemma 1], and the condition $\lim_{n \rightarrow \infty} c_n = 0$ there can be cancelled however.

Lemma 2. Let $\{c_n\}$ be any given complex O -regularly varying quasimonotone sequence. Then $\{c_n\}$ satisfies (1) for $N_0 = 1$.

The proof can be copied from Zhou and Le [11, Theorem 3] with omitting the condition $\lim_{n \rightarrow \infty} c_n = 0$ there.

⁴As usual, $A_n \approx B_n$ means that there is a positive constant $C > 0$ independent of n such that $C^{-1}B_n \leq A_n \leq CB_n$.

We obviously see that, from Lemma 1 and Lemma 2, if $\mathbf{c} = \{c_n\}_{n=1}^{\infty}$ is any given complex O -regularly varying quasimonotone sequence, then $\{c_n\} \in \text{GBV}$. On the other hand, since the converse is not true (see [11]), the class GBV gives an essential and explicit generalization to the class of O -regularly varying quasimonotone sequences.

From $\sum_{k=1}^{\infty} k^{-\alpha} \sin 2^k x =: \sum_{n=1}^{\infty} b_n \sin nx$, $\alpha > 1$, we can clearly see that, for any $\epsilon > 0$, $n^\epsilon b_n \rightarrow \infty$, $n \rightarrow \infty$, therefore, the condition (1), in general sense, cannot be further generalized.

Throughout the paper, C denotes a positive constant (which is independent of n and $x \in [0, 2\pi]$) not necessarily the same at each occurrence. In some specific cases, we also use $M(\mathbf{c})$ to indicate a positive constant only depending upon the sequence $\mathbf{c}$.

§2. Results and Proofs

We present the main result of the paper.

Theorem 1. *Let $\{\hat{f}(n)\}_{n=0}^{\infty} \in \text{GBV}$ and $\{\hat{f}(n) + \hat{f}(-n)\}_{n=1}^{\infty} \in \text{GBV}$, and*

$$f(x) = \sum_{n=-\infty}^{\infty} \hat{f}(n) e^{inx}.$$

Then $f \in C_{2\pi}$ if and only if

$$\sum_{n=1}^{\infty} |\hat{f}(n) + \hat{f}(-n)| < \infty \quad (2)$$

and

$$\lim_{n \rightarrow \infty} n \hat{f}(n) = 0. \quad (3)$$

Furthermore, if $f \in C_{2\pi}$, then

$$\begin{aligned} E_n(f) &\approx \max_{1 \leq k \leq n} k \left(|\hat{f}(n+k)| + |\hat{f}(-n-k)| \right) + \max_{k \geq 2n+1} k |\hat{f}(k) - \hat{f}(-k)| \\ &\quad + \sum_{k=2n+1}^{\infty} |\hat{f}(k) + \hat{f}(-k)|. \end{aligned}$$

First we establish several lemmas.

Lemma 3 (Xie and Zhou [8, Lemma 2]). *Let $\{\hat{f}(n)\}$ satisfy*

$$\hat{f}(n) + \hat{f}(-n) \in K(\theta_1), \quad n = 1, 2, \dots,$$

for some $\theta_1 \in [0, \pi/2)$. Then $f \in C_{2\pi}$ implies that

$$\sum_{n=1}^{\infty} |\hat{f}(n) + \hat{f}(-n)| < \infty.$$

Lemma 4. Let $\{\hat{f}(n)\}_{n=0}^{\infty} \in \text{GBV}$ and $\{\hat{f}(n) + \hat{f}(-n)\}_{n=1}^{\infty} \in \text{GBV}$. Suppose $f \in C_{2\pi}$, then

$$\max_{k \geq 1} k |\hat{f}(\pm(n+k))| = O(E_n(f)).$$

Proof. Let $t_n^*(x)$ be the trigonometric polynomials of best approximation of degree n , then from an easy equality

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} \left| e^{\pm i(n+1)x} \left(\sum_{k=0}^{N-1} e^{\pm ikx} \right)^2 \right| dx = N$$

we get

$$\left| \frac{1}{2\pi} \int_{-\pi}^{\pi} (f(x) - t_n^*(x)) e^{\pm i(n+1)x} \left(\sum_{k=0}^{N-1} e^{\pm ikx} \right)^2 dx \right| \leq N E_n(f).$$

Also, the left integral within the absolute value symbols of the above inequality equals to $\sum_{k=1}^N \left(k \hat{f}(\mp(n+k)) + (N-k) \hat{f}(\mp(n+N+k)) \right)$, then

$$\begin{aligned} N E_n(f) &\geq \left| \sum_{k=1}^N \left(k \hat{f}(\mp(n+k)) + (N-k) \hat{f}(\mp(n+N+k)) \right) \right| \\ &\geq \text{Re} \left(\sum_{k=1}^N \left(k \hat{f}(\mp(n+k)) + (N-k) \hat{f}(\mp(n+N+k)) \right) \right). \end{aligned}$$

Thus by the definition of GBV,

$$\begin{aligned} N E_n(f) &\geq \sum_{k=1}^N \left(k \text{Re} \hat{f}(n+k) + (N-k) \text{Re} \hat{f}(n+N+k) \right) \\ &\geq \sum_{k=1}^N k \text{Re} \hat{f}(n+k), \end{aligned} \quad (4)$$

and

$$2N E_n(f) \geq \sum_{k=1}^N k \left(\text{Re} \hat{f}(n+k) + \text{Re} \hat{f}(-n-k) \right). \quad (5)$$

By the same argument as (4), we also have

$$2N E_{2n}(f) \geq \sum_{k=1}^{2N} k \text{Re} \hat{f}(2n+k). \quad (4')$$

Fix sufficient large n , assume

$$\max_{N/2+jN_0 \leq k < N/2+(j+1)N_0} |\hat{f}(n+k)| = |\hat{f}(n+k_j)|, \quad j = 0, 1, \dots, [N/(2N_0)] - 1,$$

$$n + N/2 \leq n + N/2 + jN_0 \leq n + k_j < n + N/2 + (j+1)N_0 \leq n + N.$$

From condition (1), we get for $0 \leq j \leq [N/(2N_0)] - 1$,

$$\begin{aligned} |\hat{f}(n+N)| &= \left| \sum_{k=n+N}^{2n+N+2jN_0-1} \Delta \hat{f}(k) + \hat{f}(2n+N+2jN_0) \right| \\ &\leq \sum_{n+N/2+jN_0 \leq k \leq 2n+N+2jN_0} (|\Delta \hat{f}(k)| + |\hat{f}(2n+N+2jN_0)|) \\ &\leq M(\hat{f})(|\hat{f}(n+k_j)| + |\hat{f}(2n+N+2jN_0)|) \\ &\leq M(\hat{f}, \theta_1) \left(\operatorname{Re} \hat{f}(n+k_j) + \operatorname{Re} \hat{f}(2n+N+2jN_0) \right), \end{aligned}$$

therefore, together with (4) and (4'), we deduce that

$$\begin{aligned} 3NE_n(f) &\geq NE_n(f) + 2NE_{2n}(f) \geq \sum_{k=1}^N k \operatorname{Re} \hat{f}(n+k) + \sum_{k=1}^{2N} k \operatorname{Re} \hat{f}(2n+k) \\ &\geq \sum_{j=0}^{[N/(2N_0)]-1} k_j \operatorname{Re} \hat{f}(n+k_j) + \sum_{j=0}^{[N/(2N_0)]-1} (N+2jN_0) \operatorname{Re} \hat{f}(2n+N+2jN_0) \\ &\geq C \sum_{j=0}^{[N/(2N_0)]-1} k_j \left(\operatorname{Re} \hat{f}(n+k_j) + \operatorname{Re} \hat{f}(2n+N+2jN_0) \right) \\ &\geq C \sum_{j=0}^{[N/(2N_0)]-1} k_j |\hat{f}(n+N)| \geq C |\hat{f}(n+N)| \sum_{j=0}^{[N/(2N_0)]-1} (N/2 + jN_0) \\ &\geq CN_0^{-1} N^2 |\hat{f}(n+N)|. \end{aligned}$$

Finally, we achieve that, for $N \geq 1$,

$$N |\hat{f}(n+N)| \leq CE_n(f),$$

or in other words,

$$\max_{k \geq 1} k |\hat{f}(n+k)| \leq CE_n(f).$$

At the same time, starting from (5), since $\{\hat{f}(n) + \hat{f}(-n)\}_{n=1}^\infty \in \text{GBV}$, a similar argument leads to that

$$\max_{k \geq 1} k |\hat{f}(n+k) + \hat{f}(-n-k)| = O(E_n(f)).$$

Thus

$$\max_{k \geq 1} k |\hat{f}(-n-k)| \leq \max_{k \geq 1} k |\hat{f}(n+k) + \hat{f}(-n-k)| + \max_{k \geq 1} k |\hat{f}(n+k)| = O(E_n(f)).$$

Lemma 4 is completed.

As an important application, we write the following corollary of Lemma 4 as a theorem.

Theorem 2. *Let $\{\hat{f}(n)\}_{n=0}^{\infty} \in \text{GBV}$ and $\{\hat{f}(n) + \hat{f}(-n)\}_{n=1}^{\infty} \in \text{GBV}$. If $f \in C_{2\pi}$, then*

$$\lim_{n \rightarrow \infty} n |\hat{f}(\pm n)| = 0.$$

Corollary. *Let $\{b_n\}_{n=1}^{\infty} \in \text{GBV}$ be a real sequence⁵. Suppose $g(x) = \sum_{n=1}^{\infty} b_n \sin nx \in C_{2\pi}$, then*

$$\lim_{n \rightarrow \infty} n b_n = 0.$$

Lemma 5 (Xiao, Xie and Zhou [6, Lemma 4]). *Let $f \in C_{2\pi}$, $\hat{f}(n) + \hat{f}(-n) \in K(\theta_1)$, $n = 1, 2, \dots$, for some $0 \leq \theta_1 < \pi/2$. Then*

$$\sum_{k=2n+1}^{\infty} |\hat{f}(k) + \hat{f}(-k)| = O(E_n(f)).$$

Lemma 6. *Let $\{\hat{f}(n)\}_{n=0}^{\infty} \in \text{GBV}$ and $\{\hat{f}(n) + \hat{f}(-n)\}_{n=1}^{\infty} \in \text{GBV}$, then*

$$\left| \sum_{k=1}^n \hat{f}(\pm(n+k)) \sin kx \right| = O \left(\max_{1 \leq k \leq n} k \left(|\hat{f}(n+k)| + |\hat{f}(-n-k)| \right) \right)$$

holds uniformly for any $x \in [0, \pi]$.

Proof. The cases $x = 0$ and $x = \pi$ are trivial. Suppose $0 < x < \pi$. When $0 < x \leq \pi/n$, by the inequality $|\sin x| \leq |x|$ we have

$$\left| \sum_{k=1}^n \hat{f}(n+k) \sin kx \right| \leq \frac{\pi}{n} \sum_{k=1}^n k |\hat{f}(n+k)| \leq \pi \max_{1 \leq k \leq n} k |\hat{f}(n+k)|. \quad (6)$$

On the other hand, in case $\pi/n < x$, we find an natural number $m < n$ such that $m \leq \pi/x < m+1$, then

$$\left| \sum_{k=1}^n \hat{f}(n+k) \sin kx \right| \leq \frac{\pi}{m} \sum_{k=1}^m k |\hat{f}(n+k)| + \left| \sum_{k=m+1}^n \hat{f}(n+k) \sin kx \right| =: |I_1| + |I_2|.$$

It is clear that

$$|I_1| \leq \pi \max_{1 \leq k \leq m} k |\hat{f}(n+k)|. \quad (7)$$

⁵It means $b_n \geq 0$ and satisfies condition (1).

By Abel transformation,

$$I_2 = \sum_{k=m+1}^{n-N_0-1} \Delta \hat{f}(n+k) \sum_{j=1}^k \sin jx + \hat{f}(2n-N_0) \sum_{j=1}^{n-N_0} \sin jx - \hat{f}(n+m+1) \sum_{j=1}^m \sin jx \\ + \sum_{k=n-N_0+1}^n \hat{f}(n+k) \sin kx,$$

so that⁶

$$|I_2| \leq (m+1) \sum_{k=m+1}^{n-N_0-1} |\Delta \hat{f}(n+k)| + C \max_{m < k \leq n} k |\hat{f}(n+k)| + N_0 \max_{n-N_0 < k \leq n} |\hat{f}(n+k)|.$$

But by the condition of Lemma 6, taking a natural number l such that $2^l(m+1) \leq n - N_0 - 1 < 2^{l+1}(m+1)$, and setting

$$\max_{2^j(m+1) \leq k < 2^{j+1}(m+1) + N_0} |\hat{f}(n+k)| = |\hat{f}(n+k_j)|,$$

we then have⁷

$$\sum_{k=m+1}^{n-N_0-1} |\Delta \hat{f}(n+k)| \leq \sum_{j=0}^{l+1} \sum_{k=2^j(m+1)}^{2^{j+1}(m+1)-1} |\Delta \hat{f}(n+k)| \leq M(\hat{f}) \sum_{j=0}^{l+1} |\hat{f}(n+k_j)| \\ \leq C \max_{1 \leq k \leq n} k |\hat{f}(n+k)| \sum_{j=0}^{l+1} k_j^{-1} \leq C(m+1)^{-1} \max_{1 \leq k \leq n} k |\hat{f}(n+k)| \sum_{j=0}^{\infty} 2^{-j} \\ \leq C(m+1)^{-1} \max_{1 \leq k \leq n} k |\hat{f}(n+k)|.$$

Therefore, with the above estimates, we get

$$|I_2| \leq C \max_{1 < k \leq n} k |\hat{f}(n+k)|.$$

Combining with (6) and (7), we have

$$\left| \sum_{k=1}^n \hat{f}(n+k) \sin kx \right| = O \left(\max_{1 \leq k \leq n} k |\hat{f}(n+k)| \right).$$

Now write

$$\sum_{k=1}^n \hat{f}(-n-k) \sin kx = \sum_{k=1}^n \left(\hat{f}(n+k) + \hat{f}(-n-k) \right) \sin kx - \sum_{k=1}^n \hat{f}(n+k) \sin kx.$$

⁶Note that $\sum_{j=1}^k \sin jx = O(x^{-1}) = O((m+1))$ for $m \leq \pi/x < m+1$.

⁷Note that any $k_j \leq n$.

Applying the above known estimate, by noting that both $\{\hat{f}(n)\}_{n=0}^{\infty} \in \text{GBV}$ and $\{\hat{f}(n) + \hat{f}(-n)\}_{n=1}^{\infty} \in \text{GBV}$, we get

$$\begin{aligned} \left| \sum_{k=1}^n \hat{f}(-n-k) \sin kx \right| &= O \left(\max_{1 \leq k \leq n} k |\hat{f}(n+k) + \hat{f}(-n-k)| + \max_{1 \leq k \leq n} k |\hat{f}(n+k)| \right) \\ &= O \left(\max_{1 \leq k \leq n} k (|\hat{f}(n+k)| + |\hat{f}(-n-k)|) \right). \end{aligned}$$

Lemma 6 is proved.

Similarly, we can establish the following

Lemma 7. *Let $f \in C_{2\pi}$, $\{\hat{f}(n)\}_{n=0}^{\infty} \in \text{GBV}$ and $\{\hat{f}(n) + \hat{f}(-n)\}_{n=1}^{\infty} \in \text{GBV}$, then*

$$\left| \sum_{k=m}^{\infty} \hat{f}(\pm k) \sin kx \right| = O \left(\max_{k \geq m} k (|\hat{f}(k)| + |\hat{f}(-k)|) \right)$$

holds uniformly for any $x \in [0, \pi]$ and any $m \geq 1$.

Proof. Write

$$I(x) = \sum_{k=m}^{\infty} \hat{f}(k) \sin kx.$$

Noting for $x = 0$ and $x = \pi$ that $I(x) = 0$, we may restrict x within $(0, \pi)$ without loss. Take $N = [1/x]$ and set⁸

$$I(x) = \sum_{k=m}^{N-1} \hat{f}(k) \sin kx + \sum_{k=N}^{\infty} \hat{f}(k) \sin kx =: J_1(x) + J_2(x).$$

Now

$$\begin{aligned} |J_1(x)| &\leq \sum_{k=m}^{N-1} |\hat{f}(k)| |\sin kx| \leq x \sum_{k=m}^{N-1} k |\hat{f}(k)| \\ &< x(N-1)\epsilon_m \leq \epsilon_m \end{aligned}$$

follows from $N = [1/x]$, where $\epsilon_m = \max_{k \geq m} k |\hat{f}(k)|$. By Abel's transformation, similar to the proof of Lemma 6,

$$\begin{aligned} |J_2(x)| &= \left| \sum_{k=N}^{\infty} \Delta \hat{f}(k) \sum_{v=1}^k \sin vx - \hat{f}(N) \sum_{v=1}^{N-1} \sin vx \right| \\ &\leq \sum_{k=N}^{\infty} |\Delta \hat{f}(k)| \left| \sum_{v=1}^k \sin vx \right| + |\hat{f}(N)| \left| \sum_{v=1}^{N-1} \sin vx \right| \end{aligned}$$

⁸When $N \leq m$, the same argument as in estimating J_2 can be applied to deal with $I(x) = \sum_{k=m}^{\infty} \hat{f}(k) \sin kx$ directly.

$$\leq Cx^{-1} \sum_{j=0}^{\infty} \sum_{k=2^j N}^{2^{j+1}N-1} |\Delta \hat{f}(k)| + Cx^{-1} |\hat{f}(N)| \leq C\epsilon_m \sum_{j=0}^{\infty} 2^{-j} \leq C\epsilon_m.$$

Altogether, we have

$$|I(x)| \leq C\epsilon_m,$$

that is the required result for $\sum_{k=m}^{\infty} \hat{f}(k) \sin kx$. Since $\hat{f}(-k) = \hat{f}(k) + \hat{f}(-k) - \hat{f}(k)$, by the condition, we can deduce the required result for $\sum_{k=m}^{\infty} \hat{f}(-k) \sin kx$. Lemma 7 is completed.

Proof of Theorem 1. Necessity. Suppose $f \in C_{2\pi}$. From Lemma 4, (3) clearly holds, while by Lemma 5, we see

$$\sum_{k=2n+1}^{\infty} |\hat{f}(k) + \hat{f}(-k)| = O(E_n(f)),$$

thus (2) holds.

Sufficiency. It can be deduced from Zhou and Le [11, Theorem 1] with a minor revision.

Assume $f \in C_{2\pi}$ now. By Lemma 4 and Lemma 5, we see

$$\max_{k \geq 1} k |\hat{f}(\pm(n+k))| + \sum_{k=2n+1}^{\infty} |\hat{f}(k) + \hat{f}(-k)| = O(E_n(f)).$$

On the other hand, rewrite $f(x)$ as

$$\begin{aligned} f(x) &= \sum_{k=-2n}^{2n} \hat{f}(k) e^{ikx} + i \sum_{k=2n+1}^{\infty} (\hat{f}(k) - \hat{f}(-k)) \sin kx \\ &\quad + \frac{1}{2} \sum_{k=2n+1}^{\infty} (\hat{f}(k) + \hat{f}(-k)) (e^{ikx} + e^{-ikx}), \end{aligned}$$

then we have

$$\begin{aligned} E_n(f) &\leq \left\| \sum_{k=1}^n (\hat{f}(n+k) e^{i(n+k)x} + \hat{f}(-n-k) e^{-i(n+k)x}) \right. \\ &\quad \left. - \sum_{k=1}^n (\hat{f}(n+k) e^{i(n-k)x} + \hat{f}(-n-k) e^{i(-n+k)x}) \right\| \\ &\quad + \left\| \sum_{k=2n+1}^{\infty} (\hat{f}(k) - \hat{f}(-k)) \sin kx \right\| + \sum_{k=2n+1}^{\infty} |\hat{f}(k) + \hat{f}(-k)| \\ &\leq \left\| \sum_{k=1}^n (\hat{f}(n+k) \sin kx + \hat{f}(-n-k) \sin(-kx)) \right\| + \left\| \sum_{k=2n+1}^{\infty} (\hat{f}(k) - \hat{f}(-k)) \sin kx \right\| \end{aligned}$$

$$+ \sum_{k=2n+1}^{\infty} |\hat{f}(k) + \hat{f}(-k)|.$$

Applying Lemma 6 immediately yields that

$$\left\| \sum_{k=1}^n \left(\hat{f}(n+k) \sin kx + \hat{f}(-n-k) \sin(-kx) \right) \right\| \leq \max_{1 \leq k \leq n} k \left(|\hat{f}(n+k)| + |\hat{f}(-n-k)| \right),$$

while by noting that $\hat{f}(k) - \hat{f}(-k) = 2\hat{f}(k) - (\hat{f}(k) + \hat{f}(-k))$ and both $\{\hat{f}(n)\}_{n=0}^{\infty} \in \text{GBV}$ and $\{\hat{f}(n) + \hat{f}(-n)\}_{n=1}^{\infty} \in \text{GBV}$, we deduce that

$$\left\| \sum_{k=2n+1}^{\infty} \left(\hat{f}(k) - \hat{f}(-k) \right) \sin kx \right\| = O \left(2 \max_{k \geq 2n+1} k |\hat{f}(k)| + \max_{k \geq 2n+1} k |\hat{f}(k) + \hat{f}(-k)| \right)$$

by Lemma 7. Suppose $\max_{k \geq 2n+1} k |\hat{f}(k)| = m_0 |\hat{f}(m_0)|$. Take sufficient large n such that $4n - 2N_0 \geq 3n + 1$. Assume that $2n + 1 \leq m_0 \leq 4n$. Then in view of (1), by a standard technique used in the present paper, one has

$$\begin{aligned} |\hat{f}(m_0)| &\leq \sum_{k=m_0}^{4n-2N_0} |\Delta \hat{f}(k)| + |\hat{f}(4n - 2N_0 + 1)| \\ &\leq \sum_{k=2n-N_0}^{4n-2N_0} |\Delta \hat{f}(k)| + |\hat{f}(4n - 2N_0 + 1)| \\ &\leq C \left(\max_{n-N_0 \leq k \leq n} |\hat{f}(n+k)| + |\hat{f}(4n - 2N_0 + 1)| \right). \end{aligned}$$

Hence,

$$m_0 |\hat{f}(m_0)| \leq C \left(\max_{1 \leq k \leq n} k |\hat{f}(n+k)| + (4n - 2N_0 + 1) |\hat{f}(4n - 2N_0 + 1)| \right). \quad (8)$$

Meanwhile, for $m_0 \geq 4n + 1$,

$$\begin{aligned} m_0 |\hat{f}(m_0)| &\leq \frac{1}{2} m_0 |\hat{f}(m_0) - \hat{f}(-m_0)| + \frac{1}{2} m_0 |\hat{f}(m_0) + \hat{f}(-m_0)| \\ &\leq \max_{k \geq 2n+1} k |\hat{f}(k) - \hat{f}(-k)| + O \left(\sum_{k=2n+1}^{\infty} |\hat{f}(k) + \hat{f}(-k)| \right), \end{aligned}$$

where the last inequality follows from $m_0 \geq 4n + 1$ and the following calculation:
Assume that

$$\begin{aligned} \max_{(m_0+1)/2+jN_0 \leq k < (m_0+1)/2+(j+1)N_0} |\hat{f}(k) + \hat{f}(-k)| &= |\hat{f}(k_j) + \hat{f}(-k_j)|, \\ j &= 0, 1, \dots, [(m_0 + 1)/(2N_0)] - 1, \end{aligned}$$

$$2n + 1 \leq (m_0 + 1)/2 + jN_0 \leq k_j < (m_0 + 1)/2 + (j + 1)N_0 < m_0 + 1.$$

From condition (1), we get for $0 \leq j \leq [(m_0 + 1)/(2N_0)] - 1$,

$$\begin{aligned} |\hat{f}(m_0) + \hat{f}(-m_0)| &\leq \sum_{(m_0+1)/2+jN_0 \leq k \leq m_0+1+2jN_0} |\Delta(\hat{f}(k) + \hat{f}(-k))| \\ &\quad + |\hat{f}(m_0 + 1 + 2jN_0 + 1) + \hat{f}(-m_0 - 1 - 2jN_0 - 1)| \\ &\leq M(\hat{f})|\hat{f}(k_j) + \hat{f}(-k_j)| + |\hat{f}(m_0 + 1 + 2jN_0 + 1) + \hat{f}(-m_0 - 1 - 2jN_0 - 1)|, \end{aligned}$$

thus

$$m_0|\hat{f}(m_0) + \hat{f}(-m_0)| \leq CM(\hat{f}) \sum_{k=2n+1}^{\infty} |\hat{f}(k) + \hat{f}(-k)|.$$

By the same technique, for the factor appearing in (8), we also have (note $4n - 2N_0 \geq 3n + 1$)

$$(4n - 2N_0 + 1) |\hat{f}(4n - 2N_0 + 1)| \leq \max_{k \geq 2n+1} k |\hat{f}(k) - \hat{f}(-k)| + O \left(\sum_{k=2n+1}^{\infty} |\hat{f}(k) + \hat{f}(-k)| \right).$$

From the above discussion, we see that

$$\max_{k \geq 2n+1} k |\hat{f}(k)| \leq C \max_{1 \leq k \leq n} k |\hat{f}(n + k)| + \max_{k \geq 2n+1} k |\hat{f}(k) - \hat{f}(-k)| + O \left(\sum_{k=2n+1}^{\infty} |\hat{f}(k) + \hat{f}(-k)| \right)$$

holds in any case. With condition that $\hat{f}(k) + \hat{f}(-k) \in \text{GBV}$, by a similar way we can also get

$$\begin{aligned} \max_{k \geq 2n+1} k |\hat{f}(k) + \hat{f}(-k)| &\leq C \max_{1 \leq k \leq n} k (|\hat{f}(n + k)| + |\hat{f}(-n - k)|) \\ &\quad + O \left(\sum_{k=2n+1}^{\infty} |\hat{f}(k) + \hat{f}(-k)| \right). \end{aligned}$$

Combining all the above estimates, we have completed the proof of Theorem 1.

§3. An Application

Let

$$f(x) = \sum_{n=0}^{\infty} \hat{f}(n) \cos nx.$$

The following theorem is an interesting application to a hard problem in classical Fourier analysis. Interested readers could check Zhou [10] or Xie and Zhou [9, p.112], for example.

Theorem 3. Let $\{\hat{f}(n)\} \in \text{GBV}$ be a real sequence. If $f \in C_{2\pi}$ and

$$\sum_{k=n+1}^{2n} \hat{f}(k) = O\left(\max_{1 \leq k \leq n} k \hat{f}(n+k)\right),$$

then

$$\|f - S_n(f)\| = O(E_n(f)),$$

where $S_n(f)$ is the n th Fourier partial sum of f .

References

1. A. S. Belov, *On sequential estimate of best approximations and moduli of continuity by sums of trigonometric series with quasimonotone coefficients*, Matem. Zametki, 51:4(1992), 132-134, in Russian.
2. L. Leindler, *On the uniform convergence and boundedness of a certain class of sine series*, Anal. Math., 27(2001), 279-285.
3. J. R. Nurcombe, *On the uniform convergence of sine series with quasimonotone coefficients*, J. Math. Anal. Appl., 166(1992), 577-581.
4. V. B. Stanojevic, *L^1 -convergence of Fourier series with complex quasimonotone coefficients*, Proc. Amer. Math. Soc., 86(1982), 241-247.
5. V. B. Stanojevic, *L^1 -convergence of Fourier series with O -regularly varying quasimonotone coefficients*, J. Approx. Theory, 60(1990), 168-173.
6. W. Xiao, T. F. Xie and S. P. Zhou, *The best approximation rate of certain trigonometric series*, Ann. Math. Sinica, 21A(2000), 81-88, in Chinese.
7. T. F. Xie and S. P. Zhou, *L^1 -approximation of Fourier series of complex valued functions*, Proc. Royal Soc. Edinburgh, 126A(1996), 343-353.
8. T. F. Xie and S. P. Zhou, *The uniform convergence of certain trigonometric series*, J. Math. Anal. Appl., 181(1994), 171-180.
9. T. F. Xie and S. P. Zhou, *Approximation Theory of Real Functions*, Hangzhou Univ. Press, Hangzhou, 1998.
10. S. P. Zhou, *What strong monotonicity condition on Fourier coefficients can make the ratio $\|f - S_n(f)\|/E_n(f)$ bounded?*, Proc. Amer. Math. Soc., 121(1994), 779-785.
11. R. J. Le and S. P. Zhou, *A new condition for the uniform convergence of certain trigonometric series*, Acta Math. Hungar., 108(2005), 161-169.

S. P. Zhou:
Institute of Mathematics
Zhejiang Institute of Science and Technology
Hangzhou, Zhejiang 310018 China

R. J. Le:
Department of Mathematics
Ningbo University
Ningbo, Zhejiang 315211 China
and Institute of Mathematics
Zhejiang Institute of Science and Technology
Hangzhou, Zhejiang 310018 China

Keywords uniform convergence, best approximation, quasimonotone, bounded
variation